

# Caprylic Acid Supplemented in Feed Reduces Enteric *Campylobacter jejuni* Colonization in Ten-Day-Old Broiler Chickens<sup>1,2</sup>

F. Solis de los Santos,\* A. M. Donoghue,† K. Venkitanarayanan,‡ M. L. Dirain,\*  
I. Reyes-Herrera,\* P. J. Blore,\* and D. J. Donoghue\*<sup>3</sup>

\*Poultry Science Department, University of Arkansas, Fayetteville 72701; †Poultry Production and Product Safety Research Unit, Agricultural Research Service, USDA, Fayetteville, AR 72701; and ‡Department of Animal Science, University of Connecticut, Storrs 06269

**ABSTRACT** *Campylobacter* is one of the leading causes of human foodborne illness in the United States, and epidemiological evidence indicates that poultry and poultry products are a significant source of human *Campylobacter* infections. Reducing *Campylobacter* in the intestinal tract would reduce contamination of poultry products and eggs. Caprylic acid, an 8-carbon medium-chain fatty acid has been shown to be bactericidal against several pathogenic bacteria. It has, however, not been tested in the control of *Campylobacter* in chickens. Four trials were carried out to evaluate the efficacy of caprylic acid against cecal *Campylobacter jejuni* colonization in 10-d-old chicks. In the first 2 trials, day-of-hatch chicks (n = 40 per trial) were assigned to negative controls (no *Campylobacter*, no caprylic acid), positive controls (*Campylobacter*, no caprylic acid), and a low (0.7%) and a high (1.4%) dose of caprylic acid supplemented in regular chick starter feed (n = 10 chicks/treatment). Two more trials were carried out to evaluate a wider range of caprylic acid doses on

cecal *Campylobacter* counts, in which day-of-hatch chicks (n = 90 per trial) were assigned to 9 treatments: negative controls (no *Campylobacter*, no caprylic acid) and caprylic acid doses of 0 (positive controls), 0.35, 0.525, 0.7, 0.875, 1.05, 1.225, and 1.4% (n = 10 chicks/treatment). Except for the negative controls, chicks were orally gavaged with approximately  $1 \times 10^6$  cfu *Campylobacter* on d 3. On d 10, cecal contents were collected and *Campylobacter* concentrations were determined in each trial. In all 4 trials, the 0.7% dose of caprylic acid consistently reduced *Campylobacter* content counts compared with the positive control. In trials 3 and 4, doses less than 1.05% consistently reduced cecal *Campylobacter* content in both trials. At the higher doses, caprylic acid reduced feed consumption and body weight, but did not affect feed conversion when compared with the positive controls. These data suggest that low-dose supplementation with caprylic acid in feed may reduce *Campylobacter* colonization in young chickens.

**Key words:** *Campylobacter jejuni*, body weight gain, caprylic acid, broiler chick

2008 Poultry Science 87:800–804  
doi:10.3382/ps.2007-00280

## INTRODUCTION

*Campylobacter* is one of the most commonly reported bacterial causes of human foodborne illness in the United States, and epidemiological evidence indicates poultry and poultry products as significant sources of human *Campylobacter* infection (Oosterom et al., 1984; Friedman et al., 2004; CDC, 2007). Colonization of poultry by *Campylobacter* is widespread and difficult to prevent, even with

proper biosecurity measures (Wallace et al., 1998; Berrang et al., 2000; Loc Carillo et al., 2005).

Reducing *Campylobacter* in the intestinal tract would reduce contamination of poultry products. A variety of approaches for reducing the colonization of *Campylobacter* in poultry has been explored, but with varying degrees of success. These include feeding chicks or poults competitive exclusion bacteria (Mead et al., 1996; Stern et al., 2001; Mead, 2002), enzymes such as xylanase (Fernandez et al., 2000), bacteriophage (Loc Carrillo et al., 2005; Wagenaar et al., 2005), bacteriocins (Cole et al., 2006), noncultured cecal content (Mead et al., 1996; Stern et al., 2001), fructooligosaccharides and mucin-utilizing coliforms (Schoeni and Wong, 1994), organic acids (Byrd et al., 2001; Heres et al., 2004), and antibiotics (Farnell et al., 2005).

Fatty acids, especially medium-chain fatty acids, have been reported to possess antimicrobial activities against a wide range of microorganisms (Bergsson et al., 1998,

©2008 Poultry Science Association Inc.

Received July 6, 2007.

Accepted September 19, 2007.

<sup>1</sup>Mention of a trade name, proprietary product, or specific equipment does not constitute a guarantee or warranty by the USDA and does not imply its approval to the exclusion of other products that are suitable.

<sup>2</sup>Funded in part by USDA, CSREES National Integrated Food Safety Program #2006-02429 to K. Venkitanarayanan and D. J. Donoghue.

<sup>3</sup>Corresponding author: ddonogh@uark.edu

2001; Van Immerseel et al., 2004). Caprylic acid is a medium-chain fatty acid with 8 carbons and is naturally found in breast milk, bovine milk (Jensen, 2002), and coconut oil (Jensen et al., 1990; Sprong et al., 2001). It is a food-grade compound generally recognized as safe (GRAS) by the Food and Drug Administration. Caprylic acid has been shown to be effective in killing a variety of bacterial pathogens such as *Salmonella* Enteritidis in chicken cecal contents (Vasudevan et al., 2005), and *Escherichia coli*, *Staphylococcus aureus*, and *Streptococcus agalactiae* in milk (Nair et al., 2005). In preliminary studies, caprylic acid killed *Campylobacter* in vitro (data not shown). The present study was carried out to investigate whether caprylic acid supplemented in feed would reduce *Campylobacter* in the intestines of 10-d-old broiler chickens.

## MATERIALS AND METHODS

### Experimental Birds and Housing

One-day-old male broiler chicks were obtained from a commercial hatchery. These birds were placed into floor pens with dimensions of 3.06 m<sup>2</sup> (33 ft<sup>2</sup>) previously equipped with fresh litter (pine shavings) and heaters. Birds had ad libitum access to both feed and water during all trials.

### Experimental Design

Four trials were conducted in the present study. In the first 2 trials, day-of-hatch chicks (n = 40/trial) were randomly assigned to 1 of 4 treatment groups (n = 10 per treatment group). The treatments groups included negative controls (no *Campylobacter*, no caprylic acid), positive controls (*Campylobacter*, no caprylic acid), and a low (0.7%) and a high dose (1.4%) of caprylic acid (Sigma-Aldrich, St. Louis, MO) supplemented in the feed for the entire 10-d trial. To evaluate the effect of additional doses of caprylic acid on *C. jejuni* in birds, 2 follow-up trials were conducted using day-of-hatch chicks (n = 90). The chicks were assigned to 1 of 9 treatments, including negative controls (no *Campylobacter*, no caprylic acid) and 0 (positive controls), 0.35, 0.525, 0.7, 0.875, 1.05, 1.225, or 1.4% of caprylic acid supplemented in starter feed (n = 10 birds per treatment group). In all trials, the caprylic acid treatments were supplemented in feed and supplied to chicks for the entire 10-d experimental period. At 10 d of age, chicks were killed by CO<sub>2</sub> and cecal contents collected for *Campylobacter* enumeration (see below). Feed consumption was determined by subtracting the amount of feed remaining at the end of each trial from the amount given on d 1.

### Bacterial Strains and Dose

Five strains of wild-type *Campylobacter jejuni* isolated from chickens were used to colonize the birds as described previously by this laboratory (Farnell et al., 2005).

Briefly, 10- $\mu$ L loops of each frozen strain were cultured into 5 mL of *Campylobacter* enrichment broth (CEB; International Diagnostics Group plc, Lancaster, UK) and incubated for 48 h at 42°C. For the second passage, 10- $\mu$ L loops from each strain were transferred into 5 mL of fresh CEB to be incubated for 24 h at 42°C under microaerophilic conditions (5% O<sub>2</sub>, 10% CO<sub>2</sub>, and 85% N<sub>2</sub>). Following incubation, the strains were pooled in a tube and 3 mL was transferred into a glass tube to be read in a spectrophotometer to determine the concentration of bacteria present in the culture. After measuring the absorbance (optical density), the tube containing the culture was centrifuged at 3,500  $\times$  g for 10 min. The supernatant was discarded and the pellet was resuspended in an equal amount of Butterfield's phosphate diluent (BPD; 6.8% KH<sub>2</sub>PO<sub>4</sub>; 7.2 pH). The culture was serially diluted to obtain an inoculum with a desired concentration of 4  $\times$  10<sup>6</sup> cfu/mL. On d 3 of age, chicks were orally challenged individually with 250  $\mu$ L of inoculum using a 1-mL syringe connected to a sterile stainless steel cannula.

### Cecal *Campylobacter* Determination

Cecal *Campylobacter* concentrations were enumerated by the procedure of Cole et al. (2006). Briefly, ceca from each bird were transferred to a sterile plastic bag and the contents squeezed into 15-mL tubes and serially diluted (1:10) with BPD and inoculated on labeled Campy Line agar (Line, 2001) plates. The Campy Line agar (Line, 2001) plates were incubated for 48 h at 42°C under microaerophilic conditions. Direct bacterial counts were recorded and converted to colony-forming units per milliliter of the cecal content. *Campylobacter* colonies were confirmed by latex agglutination test (Panbio Inc., Columbia, MD) and further identified as *Campylobacter jejuni* using API Campy (Biomérieux, Durham, NC; ATCC 33291 strain used as control).


### Statistical Analysis

Data were analyzed by ANOVA using the GLM procedure of SAS (SAS Institute, 2002). The number of *Campylobacter* colonies was logarithmically transformed (log<sub>10</sub> cfu/mL) before analysis to achieve homogeneity of variance (Byrd et al., 2003). Treatment means were partitioned by LSMEANS analysis (SAS Institute, 2002). A probability of *P* < 0.05 was required for statistical significance.

## RESULTS

### Cecal *Campylobacter* Concentrations

The 0.7% dose of caprylic acid in the feed consistently reduced (*P* < 0.05) cecal *Campylobacter* counts in chicks compared with the positive control treatment in all 4 trials (Figure 1, Table 1). Moreover, *C. jejuni* counts recovered from the cecal contents of birds fed 0.7% caprylic acid were lower (*P* < 0.05) than that from birds fed 1.4% caprylic acid. The 1.4% caprylic acid treatment reduced



**Figure 1.** *Campylobacter jejuni* colonization of 10-d-old chicks collected from 2 separate trials ( $n = 10$  chicks/treatment per trial). A positive control (*Campylobacter* challenging without caprylic acid), 0.7%, and 1.4% doses of caprylic acid supplemented in feed for 10 consecutive days were evaluated in each trial. Chicks were orally challenged 3 d posthatch with approximately  $1 \times 10^6$  cfu/mL of a mixture of 5 *Campylobacter jejuni* isolates. The treatment groups were fed a diet containing the caprylic acid treatments; the control group (0% caprylic acid) was fed the same commercial diet without caprylic acid. <sup>a,b</sup>Columns with no common superscript differ significantly ( $P < 0.05$ ).

cecal *Campylobacter* colonization compared with the positive control group in trials 2 and 4. At this dose, however, there was no effect on cecal *Campylobacter* concentrations in trials 1 and 3 (Figure 1, Table 1). In subsequent trials (trials 3 and 4), the supplementation of chicken feed with the lower doses of caprylic acid (0.35, 0.525, 0.7, and 0.875%) reduced the counts of *Campylobacter* in 10-d-old chickens (Table 1). In these trials, the effect of caprylic acid supplementation at 1.05, 1.225 and 1.4% on *C. jejuni* carriage by birds was inconsistent. For example, although these 3 concentrations of caprylic acid had a limited effect in reducing the pathogen counts in trial 3, their effect was greater ( $P < 0.05$ ) in trial 4. *Campylobacter* was not detected in the negative control group in any of the trials.

**Table 1.** The effect of various doses of caprylic acid on cecal *Campylobacter* counts in 10-d-old chicks (mean  $\pm$  SEM)<sup>1</sup>

Caprylic acid dose, %	Trial 3	Trial 4
0	8.3 $\pm$ 0.2 <sup>a</sup>	7.6 $\pm$ 0.6 <sup>a</sup>
0.35	5.8 $\pm$ 0.6 <sup>c</sup>	2.6 $\pm$ 0.4 <sup>e</sup>
0.527	6.3 $\pm$ 0.6 <sup>bc</sup>	2.8 $\pm$ 0.5 <sup>de</sup>
0.7	6.3 $\pm$ 0.5 <sup>bc</sup>	4.5 $\pm$ 0.9 <sup>bcd</sup>
0.875	6.1 $\pm$ 0.7 <sup>c</sup>	3.6 $\pm$ 0.8 <sup>cde</sup>
1.05	7.1 $\pm$ 0.6 <sup>abc</sup>	5.3 $\pm$ 0.8 <sup>bc</sup>
1.225	7.8 $\pm$ 0.7 <sup>ab</sup>	5.6 $\pm$ 0.9 <sup>b</sup>
1.4	7.3 $\pm$ 0.7 <sup>abc</sup>	2.8 $\pm$ 0.1 <sup>de</sup>

<sup>a-e</sup>Means within columns with no common superscript differ significantly ( $P < 0.05$ ).

<sup>1</sup>All *Campylobacter* data were log<sub>10</sub> transformed for statistical analysis. In each trial, chicks were orally challenged 3 d posthatch with approximately  $1 \times 10^6$  cfu/mL of a mixture of 5 *Campylobacter jejuni* isolates ( $n = 10$  chicks/treatment per trial). Caprylic acid was fed for the entire 10-d study.

## BW, Feed Consumption, and Feed Conversion

The BW, feed consumption, and feed conversion were averaged for all trials. Feed consumption was reduced at the 1.4% dose and BW were reduced at 1.225 and 1.4% doses when compared with the positive controls (Table 2). However, the corresponding feed conversions were not affected by any of the caprylic acid doses compared with the positive controls (Table 2).

## DISCUSSION

The results of the present study demonstrate that select doses of caprylic acid in the feed can consistently reduce enteric *Campylobacter* colonization in young chickens. In 4 separate trials, the 0.7% dose of caprylic acid reduced cecal *Campylobacter* concentrations compared with the positive controls (Figure 1, Table 1). When additional doses of caprylic acid were tested in trials 3 and 4, the lower doses ( $<1.05\%$ ) of caprylic acid were more consistent than the higher doses in reducing *Campylobacter* concentrations.

The mode of action of caprylic acid on *Campylobacter* reduction in chickens is unknown, but may be due to either a direct antimicrobial effect or an indirect effect via alterations in the enteric microflora or environment. Fatty acids such as caprylic acid directly penetrate and incorporate into bacterial plasma membranes altering the membrane permeability of the bacteria (Bergsson et al., 1998, 2001). Fatty acids may also diffuse into bacterial cells in the undissociated form and dissociate within the protoplasm leading to intracellular acidification (Sun et al., 1998). A lower intracellular pH can lead to inactivation of enzymes (Viegas and Sa-Correia, 1991), and inhibition of amino acid transport (Freese et al., 1973). Caprylic acid may also conceivably reduce enteric *Campylobacter* content by altering the amounts or ratios of other enteric microflora or change some other physical characteristic of the intestine. Similar to a pro- or prebiotic, supplementation with caprylic acid could potentially alter the enteric environmental conditions making it less favorable for *Campylobacter* colonization.

At the highest dose, caprylic acid supplementation reduced feed consumption, and it could be assumed that this reduction in feed consumption resulted in these birds consuming less total caprylic acid. This may explain why the highest doses of caprylic acid were not consistent in reducing *Campylobacter* concentrations. When total caprylic acid intake was compared, however, it was apparent that even with the reduced feed consumption at higher doses, the total caprylic acid intake by birds was greater for the higher doses. For example, at the 1.4% dose, birds consumed a total of 3.7 g of caprylic acid ( $265.3 \text{ g of feed consumed} \times 1.4\% = 3.7 \text{ g}$ ; Table 2) vs. 2.0 g for the 0.7% dose. It is unclear why the higher doses of caprylic acid were less effective than the lower doses in reducing enteric *Campylobacter* counts. When caprylic



**Table 2.** The effect of various doses of caprylic acid on feed consumption, BW, and feed conversion in chicks (mean  $\pm$  SEM)<sup>1</sup>

Caprylic acid dose, %	Feed consumption	BW	Feed conversion ratio
0	318.8 $\pm$ 18.0 <sup>a</sup>	206.8 $\pm$ 7.7 <sup>a</sup>	1.58 $\pm$ 0.14 <sup>a</sup>
0.35	320.9 $\pm$ 16.1 <sup>a</sup>	196.7 $\pm$ 1.8 <sup>a</sup>	1.66 $\pm$ 0.12 <sup>a</sup>
0.525	300.6 $\pm$ 23.8 <sup>ab</sup>	195.9 $\pm$ 0.3 <sup>ab</sup>	1.57 $\pm$ 0.12 <sup>a</sup>
0.7	286.5 $\pm$ 14.9 <sup>ab</sup>	196.2 $\pm$ 8.6 <sup>ab</sup>	1.50 $\pm$ 0.13 <sup>a</sup>
0.875	289.4 $\pm$ 7.1 <sup>ab</sup>	195.1 $\pm$ 4.4 <sup>ab</sup>	1.50 $\pm$ 0.07 <sup>a</sup>
1.05	275.4 $\pm$ 6.3 <sup>ab</sup>	184.4 $\pm$ 1.1 <sup>ab</sup>	1.52 $\pm$ 0.06 <sup>a</sup>
1.225	275.4 $\pm$ 1.1 <sup>ab</sup>	174.5 $\pm$ 9.5 <sup>b</sup>	1.62 $\pm$ 0.11 <sup>a</sup>
1.4	265.3 $\pm$ 9.2 <sup>b</sup>	179.1 $\pm$ 1.1 <sup>b</sup>	1.54 $\pm$ 0.16 <sup>a</sup>

<sup>a,b</sup>Means within columns with no common superscript differ significantly ( $P < 0.05$ ).

<sup>1</sup>Data represent feed consumption, BW, and feed conversion of 10-d-old chicks averaged from separate trials. In each trial, chicks were orally challenged 3 d posthatch with approximately  $4 \times 10^5$  cfu/mL of a mixture of 5 *Campylobacter jejuni* isolates ( $n = 10$  chicks/treatment per trial). Caprylic acid was fed for the entire 10-d study.

acid was evaluated in vitro, all doses were equally effective at killing *Campylobacter* (data not shown).

The ability of caprylic acid at the highest dose to reduce feed consumption may be due to its effect on the satiety center as suggested by Cave (1982). In other words, consumption of high amounts of caprylic acid makes the bird feel full, although it has not consumed as much feed as control birds. Similar to our study results, reductions in feed consumption following caprylic acid consumption were reported by Cave (1982) and Traul et al. (2000). It is doubtful that reduced feed consumption is associated with toxicity because medium-chain fatty acids have a high degree of safety with no reported toxicity in acute toxicity tests performed orally, parenterally, or by the dermal route in several animal species, including chickens (Traul et al., 2000). These researchers reported that medium-chain triglycerides caused no risk of toxicity when ingested in a balanced diet at levels as high as 50% of dietary fat. Supplementation of Miglyl 812 (caprylic/capric acid preparation) at 16% level in the diet for 3 wk was found to be nontoxic in chickens, although the birds consumed less feed compared with control birds (Traul et al., 2000). Furthermore, caprylic acid is a food-grade chemical approved by the FDA as GRAS (FDA, 1981). Although a reduction in feed consumption and BW at the higher doses occurred in this study, the feed conversion ratio was not different (Table 2). Therefore, these birds were as efficient in feed utilization as control birds, but would take longer to reach market age. Although this would be a concern for producers, the problem could be avoided by using lower doses of caprylic acid. Moreover, the only dosing period tested in this study was 10 d and it may be possible to shorten the dosing period, thereby potentially preventing any decline in feed consumption while maintaining efficacy against *Campylobacter*. Studies are underway in our laboratory to test these possibilities.

In the present study, the dose of 0.7% caprylic acid consistently reduced cecal *Campylobacter* counts in 4 separate trials in 10-d-old broiler chickens. To our knowledge this is the first study showing that caprylic acid supplementation decreases the intestinal colonization of *Campylobacter* in broiler chickens. Follow-up studies are ongoing

to evaluate the efficacy of caprylic acid in market-aged birds.

## REFERENCES

- Bergsson, G., J. Arnfinnsson, S. M. Karlsson, Ó. Steingrímsson, and H. Thormar. 1998. In vitro inactivation of *Chlamydia trachomatis* by fatty acids and monoglycerides. *Antimicrob. Agents Chemother.* 42:2290–2294.
- Bergsson, G., J. Arnfinnsson, Ó. Steingrímsson, and H. Thormar. 2001. Killing of gram-positive cocci by fatty acids and monoglycerides. *APMIS* 109:670–678.
- Berrang, M. E., R. J. Buhr, and J. A. Cason. 2000. *Campylobacter* recovery from external and internal organs of commercial broiler carcasses prior to scalding. *Poult. Sci.* 79:286–290.
- Byrd, J. A., R. C. Anderson, T. R. Callaway, R. W. Moore, K. D. Knape, L. F. Kubena, R. L. Ziprin, and D. J. Nisbet. 2003. Effect of experimental chlorate product administration in the drinking water on *Salmonella typhimurium* contamination of broilers. *Poult. Sci.* 82:1403–1406.
- Byrd, J. A., B. M. Hargis, D. J. Caldwell, R. H. Bailey, K. L. Herron, J. L. McReynolds, R. L. Brewer, R. C. Anderson, K. M. Bischoff, T. R. Callaway, and L. F. Kubena. 2001. Effect of lactic acid administration in the drinking water during preslaughter feed withdrawal on *Salmonella* and *Campylobacter* contamination of broilers. *Poult. Sci.* 80:278–283.
- Cave, N. A. G. 1982. Effect of dietary short and medium-chain fatty acids on feed intake by chicks. *Poult. Sci.* 61:1147–1153.
- Centers for Disease Control. 2007. Preliminary FoodNet data on the incidence of infection with pathogens transmitted commonly through food—10 states, United States, 2006. *Morb. Mortal. Wkly. Rep.* 56(14):336–339.
- FDA (Food and Drug Administration). 1981. Code of Federal Regulations 21CFR 184.1025. The Office of the Federal Register National Archives and Records Administration, Washington, DC.
- Cole, K., M. B. Farnell, A. M. Donoghue, N. J. Stern, E. A. Svetoch, B. N. Eruslanov, L. I. Volodina, Y. N. Kovalev, V. V. Perelygin, E. V. Mitsevich, I. P. Mitsevich, V. P. Levchuk, V. D. Pokhilenko, V. N. Borzenkov, O. E. Svetoch, T. Y. Kudryavtseva, I. Reyes-Herrera, P. J. Blore, F. Solis de los Santos, and D. J. Donoghue. 2006. Bacteriocins reduce *Campylobacter* colonization and alter gut morphology in turkey poults. *Poult. Sci.* 85:1570–1575.
- Farnell, M. B., A. M. Donoghue, K. Cole, I. Reyes-Herrera, P. J. Blore, and D. J. Donoghue. 2005. *Campylobacter* susceptibility to ciprofloxacin and corresponding fluoroquinolone concentrations within the gastrointestinal tracts of chickens. *J. Appl. Microbiol.* 99:1043–1059.

- Fernandez, F., R. Sharma, M. Hinton, and M. R. Bedford. 2000. Diet influences the colonization of *Campylobacter jejuni* and distribution of mucin carbohydrates in the chick intestinal tract. *Cell. Mol. Life Sci.* 57:1793–1801.
- Freese, E., C. W. Sheu, and E. Galliers. 1973. Function of lipophilic acids as antimicrobial food additives. *Nature* 241:321–325.
- Friedman, C. R., R. M. Hoekstra, M. Samuel, R. Marcus, J. Bender, B. Shiferaw, S. Reddy, S. D. Ahuja, D. L. Helfrick, F. Hardnett, M. Carter, B. Anderson, and R. V. Tauxe. 2004. Risk factors for sporadic *Campylobacter* infection in the United States: A case-control study in FoodNet sites. *Clin. Infect. Dis.* 38:S285–S296.
- Heres, L., B. Engel, H. A. P. Urlings, J. A. Wagenaar, and F. van Knapen. 2004. Effect of acidified feed on susceptibility of broiler chickens to intestinal infection by *Campylobacter* and *Salmonella*. *Vet. Microbiol.* 99:259–267.
- Jensen, R. G. 2002. The composition of bovine milk lipids: January 1995 to December 2000. *J. Dairy Sci.* 85:295–350.
- Jensen, R. G., A. M. Ferris, C. J. Lammi-Keefe, and R. A. Henderson. 1990. Lipids of bovine and human milks: A comparison. *J. Dairy Sci.* 73:223–240.
- Line, J. E. 2001. Development of a selective differential agar for isolation and enumeration of *Campylobacter* spp. *J. Food Prot.* 64:1711–1715.
- Loc Carillo, C., R. J. Atterbury, A. El-Shibiny, P. L. Connerton, E. Dillon, A. Scott, and I. F. Connerton. 2005. Bacteriophage therapy to reduce *Campylobacter jejuni* colonization of broiler chickens. *Appl. Environ. Microbiol.* 71:6554–6563.
- Mead, G. C. 2002. Factors affecting intestinal colonization of poultry by *Campylobacter* and role of microflora in control. *Worlds Poult. Sci. J.* 58:169–178.
- Mead, G. C., M. J. Scott, T. J. Humphrey, and K. McAlpine. 1996. Observations on the control of *Campylobacter jejuni* infection of poultry by 'competitive exclusion'. *Avian Pathol.* 25:69–79.
- Nair, M. K., J. Joy, P. Vasudevan, L. Hinckley, T. A. Hoagland, and K. S. Venkitanarayanan. 2005. Antibacterial effect of caprylic acid and monicaprylin on major bacterial mastitis pathogens. *J. Dairy Sci.* 88:3488–3495.
- Oosterom, J., C. H. den Uyl, J. R. J. Banffer, and J. Huisman. 1984. Epidemiological investigations on *Campylobacter jejuni* in households with primary infection. *J. Hyg. (Camb.)* 92:325–332.
- SAS Institute. 2002. SAS/STAT User's Guide: Release 9.03 edition. SAS Institute Inc., Cary, NC.
- Schoeni, J. L., and A. C. L. Wong. 1994. Inhibition of *Campylobacter jejuni* colonization in chicks by defined competitive exclusion bacteria. *Appl. Environ. Microbiol.* 60:1191–1197.
- Sprong, R. C., M. F. E. Hulstein, and R. van der Meer. 2001. Bactericidal activities of milk lipids. *Antimicrob. Agents Chemother.* 45:1298–1301.
- Stern, N. J., P. Fedorka-Cray, J. S. Bailey, N. A. Cox, S. E. Craven, K. L. Hiett, M. T. Musgrove, S. Ladely, D. Cosby, and G. C. Mead. 2001. Distribution of *Campylobacter* spp. in selected U. S. poultry production and processing operations. *J. Food Prot.* 64:1705–1710.
- Sun, C. Q., C. J. O'Connor, S. J. Turner, G. D. Lewis, R. A. Stanley, and A. M. Robertson. 1998. The effect of pH on the inhibition of bacterial growth by physiological concentrations of butyric acid: Implications for neonates fed on suckled milk. *Chem. Biol. Interact.* 113:117–131.
- Traul, K. A., A. Driedger, D. L. Ingle, and D. Nakhasi. 2000. Review of the toxicologic properties of medium chain triglycerides. *Food Chem. Toxicol.* 38:79–98.
- Van Immerseel, F., J. De Buck, F. Boyen, L. Bohez, F. Pasmans, J. Volf, M. Sevcik, I. Rychlik, F. Haesebrouck, and R. Ducatelle. 2004. Medium-chain fatty acids decrease colonization and invasion through *hilA* suppression shortly after infection of chickens with *Salmonella enterica* serovar Enteritidis. *Appl. Environ. Microbiol.* 70:3582–3587.
- Vasudevan, P., M. Patrick, M. Nair, T. Annamalai, M. Darre, M. Khan, and K. Venkitanarayanan. 2005. In vitro inactivation of *Salmonella* Enteritidis in chicken cecal contents by caprylic acid. *J. Appl. Poult. Res.* 14:122–125.
- Viegas, C. A., and I. Sa-Correia. 1991. Activation of plasma membrane ATPase of *Saccharomyces cerevisiae* by octanoic acid. *J. Gen. Microbiol.* 137:645–651.
- Wagenaar, J. A., M. A. P. Van Bergen, M. A. Mueller, T. M. Wassenaar, and R. M. Carlton. 2005. Phage therapy reduces *Campylobacter jejuni* colonization in broilers. *Vet. Microbiol.* 109:275–283.
- Wallace, J. S., K. N. Stanley, and K. Jones. 1998. The colonization of turkeys by thermophilic *Campylobacters*. *J. Appl. Microbiol.* 85:224–230.